

# A New Species of *Chromodoris*

(Opisthobranchia : Nudibranchia)

## from Tropical West America

BY

HANS BERTSCH

Donner Laboratory and Department of Zoology, University of California, Berkeley, California 94720

(1 Plate; 2 Text figures)

DURING THE PAST few years, Dr. Antonio J. Ferreira and Mr. and Mrs. Roy Poorman have collected 4 specimens of a small chromodorid from tropical west American waters. Preserved specimens and color transparencies of the living animals were given to me for examination. A careful comparison of taxonomically important anatomical features has shown these specimens to be members of a new species.

### HOLOHEPATICA

#### Doridoidea

#### CHROMODORIDIDAE

#### Chromodoridinae

#### *Chromodoris antonii* Bertsch, spec. nov.

**Material Examined:** 1) **Holotype:** One specimen (4 mm long preserved) collected in about 9 m of water, at Punta Aguja (near Mulege), Baja California, Mexico (approximately 26°53' N; 111°56' W), on July 28, 1974, by Antonio J. Ferreira. This dissected specimen and its mounted radula have been deposited in the collections of the Los Angeles County Natural History Museum, LACM Type Series, No. 1810.

2) **Paratypes:** One specimen (4.5 mm long preserved) collected subtidally in about 10 - 12 m of water, Isla San Jose, Baja California, Mexico (approximately 24°53' N; 110°34' W), on August 14, 1973, by A. J. Ferreira. The dissected specimen and its mounted radula are in the LACM Type Series, No. 1811.

3) One specimen (4 mm long preserved), collected subtidally in about 28 m of water, Los Islotes, Baja California, Mexico (24°36' N; 110°24' W) on August 14, 1973, by A. J. Ferreira. The intact specimen is also in the LACM Type Series, No. 1812.

4) One specimen (alive, 10 mm long) collected in Santiago Bay, near Manzanillo, Colima, Mexico (approximately 19°06' N; 104°23' W), in January, 1976, by Laura Shy and Forrest & Roy Poorman while diving. The dissected specimen and its mounted radula are in the LACM Type Series, No. 1813.

This species has been collected only from subtidal regions, offshore along the Baja California coast of the central and southern Gulf of California, and from the southwestern coast of mainland Mexico. The known range of *Chromodoris antonii* extends some 1260 km in a north-west - southwest direction in the tropical west American marine faunal province.

**Description:** Lengths of preserved specimens vary from 3 to 4.5 mm; specimen 4 was measured alive as 10 mm long. Mantle margin extends past the sides of the animal's body, with the tail protruding out behind the posterior mantle margin about 1/9 the distance of the animal's total length (Figure 1).

Coloration consists basically of shades of blue, magenta, black, yellow-orange and white. A complete yellow-orange line encircles the rim of the notum; a black line immediately borders the entire inner side of the yellow-orange band. A wide area of light blue covers the rest of the lateral notal region. This zone is divided by a blackish line concentric with the 2 outermost bands. The central dorsal region (from between the rhinophores to the anterior and lateral sides of the gills) bears a light magenta

Table 1

Radular sizes and teeth counts of *Chromodoris antonii*

| Specimen   | Length<br>mm | Width<br>mm | W:L<br>Ratio | Rows of<br>Teeth | Number of Teeth<br>Per Half-Row |
|------------|--------------|-------------|--------------|------------------|---------------------------------|
| Material 1 | 0.61         | 0.3         | 1:2.03       | ~56              | ~35                             |
| Material 2 | 0.75         | 0.27        | 1:2.78       | ~78              | ~41                             |
| Material 4 | 0.64         | 0.31        | 1:2.06       | ~54              | ~32                             |
| Average    | 0.67         | 0.29        | 1:2.31       | ~63              | ~36                             |

color, with darker splotches scattered randomly. Running lengthwise down the center of the notum is a thick, broken white line, bordered by a slight yellowish tinge. Rhino-



Figure 1

*Chromodoris antonii* Bertsch, spec. nov.

Drawing of living animal (from a color transparency  
by A. J. Ferreira). Holotype (LACM 1810)

phores are light magenta proximally, and black the distal  $\frac{1}{2}$  of their length. The 6 or 7 pinnate gill plumes are pinkish-white basally, each tipped with black. Rim of branchial pouch is tinged yellowish. The dorsal portion of the foot, extending past the mantle, is rimmed by black, with successive light blue and dark blue regions, with a small broken white line along its center.

The approximate radular sizes and formulae of specimens 1, 2, and 4 are given in Table 1. Because of their extremely small size, the radulae were damaged in mounting; exact measurements, and tooth and row counts of the radulae were not possible. The radula has no rachidian tooth. The lateral teeth bear long denticles, approximately  $\frac{1}{2}$  the total width of the erect hook along an antero-posterior axis (Figure 2). There is no prominent cusp thicker and longer than the succeeding denticles. Instead, this structure is reduced to approximately the same size and thickness as the denticles, and it is often actually shorter than the immediately adjacent denticle. In the following description, this "cusp" is counted simply as a denticle. The inner teeth of each half-row have 4 denticles (Figures 3, 4, 5). The innermost tooth does not appear to have additional denticles on its inner face. In the central portion of each half-row the teeth have 6 - 7

(sometimes 8) denticles (Figures 6 and 7). The outermost lateral teeth have only 4 or 5 denticles (Figure 8). All the teeth have a base not clearly set off from the erect denticulate shaft; the postero-dorsal surface of the tooth curves evenly upwards from the posterior portion of the base (Figure 5); outer teeth have a very short base (Figure 2).



Figure 2

*Chromodoris antonii* Bertsch, spec. nov.

Outer lateral teeth from specimen 2  
(LACM 1811) approximately  $\times 1500$

**Discussion:** Five genera of Chromodorididae have been reported from the Pacific coast of North America. Species of the genus *Cadlina* are differentiated from *Chromodoris antonii* because they 1) have a rachidian tooth with pronounced denticulations on the erect shaft; 2) are usually colored whitish or yellow or shades of both; and 3) ecologically are predominantly members of temperate, cooler water faunal provinces.

*Chromodoris antonii* can be separated from species of the remaining tropical chromodorid genera on the basis of coloration (cf. RUDMAN, 1973, for a discussion about the species-specificity of coloration among chromodorids) and radular characteristics.

*Chromodoris banksi* Farmer, 1963, *C. baumanni* Bertsch, 1970, *C. marislac* Bertsch, in Bertsch *et al.*, 1973, *C. norrisi* Farmer, 1963, and *C. sedna* (Marcus & Marcus, 1967) all have a white notal background color and a prominent cusp on the shaft of each lateral tooth. *Chromodoris mc-*



*farlandi* Cockerell, 1901, is violet, with a yellow-orange band encircling the notum edge, and 3 longitudinal yellowish lines in the mid-lateral and central portions of the dorsum; its teeth have pronounced cusps. The teeth of *C. porterae* Cockerell, 1901, resemble *C. antonii* in overall morphology; however, the coloration (blue with a yellow-orange line surrounding the central portion of the dorsum) of *C. porterae* is distinctly different from *C. antonii*. *Chromodoris dalli* Bergh, 1879a, is an enigmatic species, but its teeth have the typical prominent cusp. *Chromodoris glauca* Bergh, 1879b, is a synonym of *Hypselodoris californiensis* (Bergh, 1879a) (BERTSCH, in prep.).

*Thorunna tura* (Marcus & Marcus, 1967) and *T. lapislazuli* Bertsch & Ferreira, 1974, have numerous yellow-orange dots on the notum and lack either a peripheral orange or black line on the notal edge. Moreover, both species have a much broader based innermost lateral tooth; *T. tura* has a higher ratio of teeth per half-row : total rows of teeth than does *Chromodoris antonii*, and *T. lapislazuli* has bicuspid outer lateral teeth.

*Hypselodoris* sp. (figured in MARCUS & MARCUS, 1967: 177, fig. 30), *H. agassizii* (Bergh, 1894), and *H. californiensis* all have bicuspid teeth with large distal cusps, and a dark blue body color with yellow dots or streaks on the dorsum *Hypselodoris aegialia* (Bergh, 1904) has bicuspid teeth, without additional denticles.

*Felimida sphoni* Marcus, 1971, has a "red cross" coloration pattern; a distinct, large cusp on each tooth; and denticles on both the inner and outer faces of the 4 innermost lateral teeth.

The specific name *antonii* is chosen to honor Dr. Antonio J. Ferreira, my esteemed friend and fellow worker, who graciously has shared with me his time, interest, and enthusiasm, and has contributed significantly to the knowledge of polyplacophoran and opisthobranch molluscan taxonomy.

## ACKNOWLEDGMENTS

I thank Dr. Antonio J. Ferreira for giving me the specimens he collected; Mr. Gale G. Sphon, Los Angeles County Natural History Museum, for lending the specimen

collected by Mr. and Mrs. Poorman; Dr. Thomas L. Hayes, Donner Laboratory, University of California, Berkeley, for working time on the scanning electron microscope; Dr. Richard Strohman, Chairperson, Department of Zoology, University of California, Berkeley, for providing research funds; and Ms. Susan Pohlman and Mr. Robert Stevens for assistance with the text figures.

## Literature Cited

- BERGH, LUDWIG SOPHUS RUDOLF  
1879a. On the nudibranchiate gastropod Mollusca of the North Pacific Ocean, with special reference to those of Alaska. Proc. Acad. Nat. Sci. Philadelphia, prt. 1; 31: 127-188; pls 1-8 (10 May 1879)  
1879b. Neue Chromodoriden. Malakoz. Blätter, N. F. 1: 87-116; plt. 3  
1894. Reports on the dredging operations off the west coast of Central America to the Galapagos, to the west coast of Mexico, and in the Gulf of California, in charge of Alexander Agassiz, carried on by the U. S. Fish Commission steamer "Albatross," during 1891, Lieut. Commander Z. L. Tanner, U. S. N., commanding. XIII. Die Opisthobranchien. Bull. Mus. Comp. Zool. Harvard Univ. 25 (10): 125 to 233; pls. 1-12 (October 1894)  
1904. Malacologische Untersuchungen: Nudibranchiata. In: C. SEMPER, Reisen im Archipel der Philippinen, Wissenschaftliche Resultate 9 (6): (1): 1-55; pls. 1-4 (2 February 1904)
- BERTSCH, HANS  
1970. Opisthobranchs from Isla San Francisco, Gulf of California, with the description of a new species. Contrib. Sci., Santa Barbara Mus. Nat. Hist. 2: 1-16; 13 text figs. (1 December 1970)
- BERTSCH, HANS & ANTONIO J. FERREIRA  
1974. Four new species of nudibranchs from tropical west America. The Veliger 16 (4): 343-353; 7 pls. (1 April 1974)
- BERTSCH, HANS, ANTONIO J. FERREIRA, WESLEY M. FARMER & THOMAS L. HAYES  
1973. The genera *Chromodoris* and *Felimida* (Nudibranchia: Chromodorididae) in tropical west America: distributional data, description of a new species, and scanning electron microscopic studies of radulae. The Veliger 15 (4): 287-294; 3 pls.; 3 text figs. (1 April 1973)
- COCKERELL, THEODORE DRU ALISON  
1901. Pigments of nudibranchiate Mollusca. Nature 65 (1674): 79-80 (28 November 1901)
- FARMER, WESLEY MERRILL  
1963. Two new opisthobranch mollusks from Baja California. Trans. San Diego Soc. Nat. Hist. 13 (6) 81-84; 1 plt.; 1 text fig. (27 September 1963)
- MARCUS, EVELINE DU BOIS REYMOND  
1971. On some euthyneuran gastropods from the Indian and Pacific Oceans. Proc. Malacol. Soc. London 39 (5): 355-369; 20 text figs. (August 1971)
- MARCUS, EVELINE DU BOIS-REYMOND & ERNST MARCUS  
1967. American opisthobranch mollusks. Studies in tropical oceanography (Univ. Miami Inst. Marine Sci., Miami, Florida), no. 6: viii + 256 pp.; figs. 1-155 + 1-95 (22 December 1967)
- RUDMAN, W. B.  
1973. Chromodorid opisthobranch Mollusca from the Indo-West Pacific. Zool. Journ. Linn. Soc. 52 (3): 175-199; 2 pls.; 7 text figs. (4 July 1973)

## Explanation of Figures 3 to 8

Scanning Electron Micrographs of *Chromodoris antonii* Radulae

Figure 3: Inner lateral teeth, holotype (LACM 1810) ca.  $\times 570$

Figure 4: Inner and central lateral teeth, specimen 4

(LACM 1813) ca.  $\times 570$

Figure 5: Inner lateral teeth, holotype (LACM 1810) ca.  $\times 1700$

Figure 6: Central lateral teeth (near middle of half-row), specimen 2 (LACM 1811) ca.  $\times 1700$

Figure 7: Outer lateral teeth, specimen 4 (LACM 1813) ca.  $\times 1700$

Figure 8: Outermost lateral teeth, specimen 4 (LACM 1813) ca.  $\times 1700$